

University of Groningen

Collateral Ventilation

Slebos, Dirk-Jan; Shah, Pallav L.

Published in:
Respiration

DOI:
[10.1159/000456672](https://doi.org/10.1159/000456672)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2017

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Slebos, D.-J., & Shah, P. L. (2017). Collateral Ventilation: Friend or Foe in Patients with Severe Emphysema. *Respiration*, 93(4), 232-233. <https://doi.org/10.1159/000456672>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Collateral Ventilation: Friend or Foe in Patients with Severe Emphysema

Dirk-Jan Slebos^a Pallav L. Shah^{b, c}

^aDepartment of Pulmonary Diseases, University of Groningen, University Medical Center Groningen, Groningen, The Netherlands; ^bThe National Institute for Health Research Unit, Royal Brompton and Harefield NHS Foundation Trust and Imperial College, and ^cChelsea and Westminster Hospital NHS Foundation Trust, London, UK

Over the past decade, significant improvements have been made in developing nonpharmacological interventions to enhance lung mechanics in patients with severe emphysema [1]. Collateral ventilation has been an influential factor for some of the therapies and is defined as ventilation of the alveoli via pathways that bypass the normal bronchial airways. The initial randomized controlled trials with endobronchial valves failed to demonstrate clinically meaningful results as a proportion of the treated patients had collateral ventilation and hence did not develop lobar atelectasis [2]. Once the role of collateral ventilation was appreciated and patients were more carefully selected on the basis of formal measurements, remarkable benefits have been observed with mean improvements in FEV₁ between 20 and 25% [3–5]. Such is the importance of collateral ventilation that surrogate markers for collateral ventilation such as fissure integrity are formally assessed on the computed tomography scans at the screening stage and in the majority of patients formal measurements of collateral ventilation are performed by the Chartis procedure as recommended by the latest Best Practice Guidelines [6].

A significant proportion of patients with emphysema have collateral ventilation and hence alternative bronchoscopic techniques have been developed that work in-

dependently of this physiological phenomenon. For this patient group, both endobronchial coils [7, 8] as well as sclerosing therapies [9–11] have shown potential as treatment options. Collateral ventilation thus seems the most important denominator in both chronic obstructive pulmonary disease and in bronchoscopic lung volume reduction. Where there is a dominant chronic obstructive pulmonary disease component, there is mucosal inflammation, bronchospasm, irregular airways, and mucus plugging. In this situation, the resistance of the airways is usually high leading to progressive air trapping, hypoxemia and hypercapnia. With progressive emphysema, there is destruction of the alveolar parenchyma increasing both intra- and interlobar collateral ventilation. Very early studies in excised emphysematous lungs confirmed that the resistance to collateral flow is much lower than the resistance in the airways [12]. This allows the exchange of gas even in obstructed segments and may account for the relative preservation of hypoxemia in emphysema patients.

In the past, several attempts have been made to take advantage of the abundant presence of collateral ventilation in severe emphysema by creating fenestrations – a so-called “airway bypass” – between the areas of trapped air and places where this air can easily be released. Early

work in this field both showed potential using a transthoracic airway bypass approach [13, 14], as well as creating a bronchoscopic transbronchial airway bypass [15].

To prove the safety and efficacy of creating extra-anatomic airway bypasses between the most diseased areas of the lung and the large airways which were supported by paclitaxel drug-eluting stents to ensure patency, a randomized, full sham-controlled, multicenter trial (the EASE trial) was performed in patients with severe homogeneous emphysema [16]. The EASE trial showed that the airway bypass improved patient outcomes directly after the procedure; however, the durability of the effects was limited, and outcomes for both the sham-controlled and treatment groups were similar long after the procedure. Although the concept of creating an airway bypass was proven in this trial, the loss of airway bypass patency due to several factors caused the loss of the observed initial benefit [16].

In this issue of *Respiration*, Snell et al. [17] report on a unique treatment in a unique patient population whom

they were able to follow for years after treatment with a transthoracic airway bypass. Creating a transthoracic airway bypass comes with a number of potential issues, such as thoracic surgery to create the bypass, with initially significant subcutaneous emphysema, frequent change of the bypass tubing, frequent interventions to clear out the bypass channel and of course also local esthetical issues. However, this paper again shows the great potential of creating functional collateral channels allowing emphysematous lungs to more easily empty and by those means improve their dyspnea symptoms. Based on this long-term follow-up, earlier reports, and our own observations [16, 18], we can only encourage further innovation and development of – preferably endoscopic – techniques that will allow this therapeutic approach to become more widely applicable for our highly symptomatic severe emphysema patients. Especially homogeneous emphysema patients with a high level of tissue destruction and who do have interlobar collateral ventilation seem to be the target patients for this therapy.

References

- Shah PL, Herth FJ, van Geffen WH, Deslee G, Slebos DJ: Lung volume reduction for emphysema. *Lancet Respir Med* 2016, Epub ahead of print.
- Sciruba FC, Ernst A, Herth FJ, Strange C, Criner GJ, Marquette CH, et al: A randomized study of endobronchial valves for advanced emphysema. *N Engl J Med* 2010;363:1233–1244.
- Klooster K, ten Hacken NH, Hartman JE, Kerstjens HA, van Rikxoort EM, Slebos DJ: Endobronchial valves for emphysema without interlobar collateral ventilation. *N Engl J Med* 2015;373:2325–2335.
- Davey C, Zoumot Z, Jordan S, McNulty WH, Carr DH, Hind MD, et al: Bronchoscopic lung volume reduction with endobronchial valves for patients with heterogeneous emphysema and intact interlobar fissures (the BeLieVeR-HiFi study): a randomised controlled trial. *Lancet* 2015;386:1066–1073.
- Koster TD, Slebos DJ: The fissure: interlobar collateral ventilation and implications for endoscopic therapy in emphysema. *Int J Chron Obstruct Pulmon Dis* 2016;11:765–773.
- Slebos DJ, Shah PL, Herth FJ, Valipour A: Endobronchial valves for endoscopic lung volume reduction: best practice recommendations from expert panel on endoscopic lung volume reduction. *Respiration* 2016, Epub ahead of print.
- Sciruba FC, Criner GJ, Strange C, Shah PL, Michaud G, Connolly TA, et al: Effect of endobronchial coils versus usual care on exercise tolerance in patients with severe emphysema: the RENEW randomized clinical trial. *JAMA* 2016;315:2178–2189.
- Slebos DJ, Hartman JE, Klooster K, Blaas S, Deslee G, Gesierich W, et al: Bronchoscopic coil treatment for patients with severe emphysema: a meta-analysis. *Respiration* 2015;90:136–145.
- Come CE, Kramer MR, Dransfield MT, Abu-Hijleh M, Berkowitz D, Bezzi M, et al: A randomised trial of lung sealant versus medical therapy for advanced emphysema. *Eur Respir J* 2015;46:651–662.
- Shah PL, Gompelmann D, Valipour A, McNulty WH, Eberhardt R, Grah C, et al: Thermal vapour ablation to reduce segmental volume in patients with severe emphysema: STEP-UP 12 month results. *Lancet Respir Med* 2016;4:e44–e45.
- Gompelmann D, Eberhardt R, Schuhmann M, Valipour A, Shah PL, Herth FJ, Kontogianni K: Lung volume reduction with vapor ablation in the presence of incomplete fissures: 12-month results from the STEP-UP randomized controlled study. *Respiration* 2016;92:397–403.
- Hogg JC, Macklem PT, Thurlbeck WM: The resistance of collateral channels in excised human lungs. *J Clin Invest* 1969;48:421–431.
- Junior RS, Neto VD, Botter M, Stirbulov R, Rivaben JH, Gonçalves R: Therapeutic application of collateral ventilation with pulmonary drainage in the treatment of diffuse emphysema: report of the first three cases. *J Bras Pneumol* 2009;35:14–19.
- Moore AJ, Cetti E, Haj-Yahia S, Carby M, Björling G, Karlsson S, Shah P, Goldstraw P, Moxham J, Jordan S, Polkey MI: Unilateral extrapulmonary airway bypass in advanced emphysema. *Ann Thorac Surg* 2010;89:899–906.
- Choong CK, Macklem PT, Pierce JA, Das N, Lutey BA, Martinez CO, Cooper JD: Airway bypass improves the mechanical properties of explanted emphysematous lungs. *Am J Respir Crit Care Med* 2008;178:902–905.
- Shah PL, Slebos DJ, Cardoso PF, Cetti E, Voelker K, Levine B, Russell ME, Goldin J, Brown M, Cooper JD, Sybrecht GW; EASE trial study group: Bronchoscopic lung-volume reduction with Exhale airway stents for emphysema (EASE trial): randomised, sham-controlled, multicentre trial. *Lancet* 2011;378:997–1005.
- Snell GI, Holsworth L, Khorramnia S, Westall GP, Williams TJ, Marasco S, Gooi JH: Feasibility and safety of a transthoracic pneumostoma airway bypass in severe emphysema patients. *Respiration* 2017;93:236–246.
- Slebos DJ, Klooster K, Erasmus M: Emphysema! *Am J Respir Crit Care Med* 2012;186:197.